

# 2021 Grape quick reference guide



Performance with more power.



Cluster	Product (Group)	PCP#	Crop (crop group)	Application placement	Formulation type	WALES mixing order	Maximum applications	Application rate*	Adjuvant	Application interval (days)	REI	PHI (days)	Control of	Comments
<b>Fungicide</b>														
Powdery mildew	<b>Aprovia Top</b> (Group 3 + 7)	31526	Grapes (13-07F)	Foliar	EC	E	4	0.26 L/acre (0.64 L/ha)	None	11 to 21	10 days to perform girdling and turning, 2 days for all other activities	21	Powdery mildew	
Powdery mildew and Botrytis	<b>Inspire Super</b> (Group 3 + 9)	30827	Grapes	Foliar	EW	E	2	0.34 to 0.6 L/acre (0.84 to 1.48 L/ha)	NIS 0.1 %v/v	15 to 21	15 days to perform girdling and turning, 7 days for all other activities	14	Anthracnose, Alternaria rot, powdery mildew, Botrytis bunch rot, black rot	Application rate differs by target.
	<b>Miravis Prime</b> (Group 7 + 12)	33207	Grapes (13-07F)	Foliar	SC	L	2	0.32 to 0.4 L/acre (0.8 to 1.0 L/ha)	None	21	1 day to perform girdling and turning, 12 hours for all other activities	14	Botrytis grey mould, powdery mildew	Powdery mildew control when applied at the timing for Botrytis grey mould.
Botrytis	<b>Switch 62.5WG</b> (Group 9 + 12)	28189	Grapes (13-07F)	Foliar	WG	W	2	314 to 395 g/acre (775 to 975 g/ha)	None	21	7 days to perform girdling and turning, 2 days for training, thinning, hand pruning, tying and leaf pulling activities, 12 hours for all other activities	7	Botrytis grey mould	
Downy mildew	<b>Revus</b> (Group 40)	29074	Grapes	Foliar	SC	L	4	0.2 L/acre (0.5 L/ha)	NIS 0.25 %v/v	7 to 10	12 hours	14	Downy mildew	
	<b>Ridomil Gold MZ</b> (Group M + 4)	28893	Grapes	Foliar	WG	W	1	1 kg/acre (2.5 kg/ha)	None	NA	24 hours	66	Downy mildew	

REI = Re-entry interval. PHI = Pre-harvest interval.

Continued on reverse

\*For complete information regarding rates, see product label. Rate information is based on product labels. In the case of ranges, the most commonly used rates are listed.

Always refer to the label to confirm application information, as well as buffer zones, PPE, etc., before application.

For more information, contact our Customer Interaction Centre at 1-877-SYNGENTA (1-877-964-3682) or visit Syngenta.ca.

Always read and follow label directions. Agri-Mek®, Aprovia®, Inspire Super®, Miravis®, Revus®, Ridomil Gold®, Switch®, Venture® and the Syngenta logo are trademarks of a Syngenta Group Company. © 2020 Syngenta

For more information, visit [Syngenta.ca/Miravis-Prime](http://Syngenta.ca/Miravis-Prime)



# 2021 Grape quick reference guide

**Miravis<sup>®</sup>Prime**  
Performance with more power.



Cluster	Product (Group)	PCP#	Crop (crop group)	Application placement	Formulation type	WALES mixing order	Maximum applications	Application rate*	Adjuvant	Application interval (days)	REI	PHI (days)	Control of	Comments
<b>Insecticide</b>														
Mites	<b>Agri-Mek<sup>®</sup>SC</b> (Group 6)	31607	Grapes	Foliar	SC	L	2	53 to 107 mL/acre (130 to 265 mL/ha)	NIS 0.1 to 0.5 %v/v	21		28	Two-spotted spider mite, European red mite	
<b>Herbicide</b>														
Grasses	<b>Venture<sup>®</sup>L</b> (Group 1)	21209	Grapes	Foliar	EC	E	1	0.24 to 0.81 L/acre (0.6 to 2 L/ha)	None	NA	12 hours	30	Annual and perennial grass weeds	Application rate differs by target.

\*For complete information regarding rates, see product label. Rate information is based on product labels.

REI = Re-entry interval. PHI = Pre-harvest interval.

## Application timing for fungicides and herbicides



Bud break	Shoot development	Bloom	Post-bloom/ Early fruit set	Berry touch/ Cluster closure	Veraison	Harvest

For more information, visit [Syngenta.ca/Miravis-Prime](http://Syngenta.ca/Miravis-Prime)